

Ocimum Species: Ethnomedicinal Uses, Phytochemistry and Pharmacological Importance

Chinedu Enegide

Department of Pharmacology, Novena University, Ogume, Delta State, Nigeria

Ofili Charles C

Department of Public and Community Health, Novena University, Ogume, Delta State, Nigeria

Plant species belonging to the *Ocimum* genus are among the most popular medicinal plants and are being used for several purposes in ethnomedicine. *Ocimum* is universally cultivated however; the geographical distribution highlights three major centers of *Ocimum* diversity, these are: the tropical areas of America, the tropical region of Asia and the tropical and subtropical regions of Africa. *Ocimum* species have been employed traditionally for several medicinal purposes including antioxidant, antibacterial, hypoglycaemic, hepatoprotective, antiviral and other medicinal purposes. Though scientists have carried-out studies and chronicled the pharmacological potentials of *Ocimum* species, documents containing these data seem to be disassembled, making it difficult to really distinguish the pharmacologically useful species from those that are not and also to get comprehensive information about the medicinal species in this genus. Hence, the aim of this review is to outline classified information on the ethnopharmacology, phytochemistry and pharmacological importance of some *Ocimum* species that have been reported by different researchers. In carrying-out this review, the search for literature was done via relevant databases including PubMed, Springer, Web of Science, Science Direct, Embase, SciFinder, Google Scholar and Scopus. The species reviewed are the most widely used from the *Ocimum* genus in traditional medicine and they are also the most researched for intended use in conventional therapeutic practice. Literature reveals that these species contain several compounds which are responsible for the numerous pharmacological activities elicited by them including antimalarial, anticancer, antidiabetic, antiinflammatory and antioxidant effects.

Introduction

In the past few years, scientists and researchers have largely focused on natural products in search of potent and safer agents for curative, prophylactic and other medicinal purposes.^[1] This have led to extensive scientific studies on suspected medicinal plants and other natural products.^[2] Scientific chronicles have shown that remarkable success have evolved from this trend, as several pharmacological agents currently in use were developed from natural products (including plants).^[3]

Plant species belonging to the *Ocimum* genus are among the most popular medicinal plants and are being used for several purposes in ethnomedicine. Of the entire genus stemming from the subfamily Nepetoideae (under the family Lamiaceae), *Ocimum* (also known as Basil) is referred to as the most the most important of them.^[4] The name *Ocimum* was derived from the word "Ozo" (Greek), which means smell.^[5] Due to its enormous use in both traditional medicine and pharmaceutical industry, and as well for perfumery purpose, it is also referred to as "king of herbs."^[6] According to the report documented by Pushpangadan and Bradu, the genus *Ocimum* with over 160 species is said to be the largest genera belonging to Lamiaceae Family.^[7] About 65 of the species are said to be native to *Ocimum*, while the others are considered as synonyms.^[1]

Ocimum is universally cultivated however, the geographical distribution highlights three major centers of Ocimum diversity. These are: the tropical areas of America, the tropical region of Asia and the tropical and subtropical regions of Africa. The African tropical rain forest has the highest number of Ocimum species.^[8] The main species of Ocimum with documented pharmacological activity includes *Ocimum gratissimum*, *Ocimum basilicum*, *Ocimum sanctum*, *Ocimum americanum*, and *Ocimum Kilimandscharicum*. These species have been reported traditionally to possess different culinary and medicinal activities. In different parts of the world, people employ plant species from this genus for several medicinal purposes such antioxidant, antibacterial, hypoglycaemic, hepatoprotective, antiviral and several other medicinal purposes.^[9-12] Though scientists have carried-out studies and chronicled the pharmacological potentials of Ocimum, documents containing these data seem to be disassembled, making it difficult to really distinguish the pharmacologically useful species from those that are not and also to get comprehensive information about the medicinal species in this genus. Hence, the aim of this review is to outline classified information on the ethnopharmacology, phytochemistry and pharmacological importance of some Ocimum species that have been reported by different researchers.

In carrying-out this review, the search for literature was done via relevant databases including PubMed, Springer, Web of Science, Science Direct, Embase, SciFinder, Google Scholar and Scopus. The keywords used for the search includes: Ocimum, Ocimum species, phytochemical, phytochemistry, pharmacological activity, pharmacological evaluation of extracts, fractions, or isolated compounds from Ocimum. Selection and inclusion of articles for the study was based on publications done in English language only. All selected manuscripts were analyzed for relevance to the topic, reported plant species, isolated chemical compounds, and evaluated biological activities.

Ethnopharmacology of Ocimum species

Literature shows that traditionally, plant species belonging to the Ocimum genus are widely employed for the treatment and management of several ailments including mental illness, diarrhea, measles, gonorrhea, rheumatism, paralysis, high fever, influenza, epilepsy, abdominal pains, cold and cough. They are also used as antipyretic, antihelmentic, antiemetic and antimalarial agents traditionally.^[13-14] The specific documented tradomedical application of the different species belonging to the Ocimum genus is summarized in table 1 below.

Species	Region	Parts	Traditional Uses	References
<i>Ocimum americanum</i>	China, India, Thailand, Nigeria, Cameroon, Mali, Guinea	Leaves	It is used for treating coughs, bronchial catarrh, ulcers, haemorrhoids, tuberculosis, stomach pains ear and eye ailments. It is also used for lowering high blood pressure, and to treat constipation, stomach ache, diabetes, diarrhea, dysentery and haemorrhoids.	[15-19]
<i>Ocimum basilicum</i>	Cameroon, Egypt, Nigeria, Guinea, Mali, Rwanda	Leaves	Used for prophylaxis and treating of cardiovascular disorders, diabetics, aches and pains, cough, headache, kidney malfunctions and diarrhea. It also used in the treatment of skin infections, snake bites and insect stings, and as a sedative.	[20-23]

<i>Ocimum gratissimum</i>	India, China, Nigeria, New Zealand, Australia	Leaves, stem, root and flowers	It is used as an antidiabetic, antiseptic, antidiarrhoeal, antitussive, antihelmintic, antipyretic, anti-inflammatory, antispasmodic and antimicrobial agent. Also used for the treatment and management of various stomach and kidney ailments, upper respiratory tract infection, pneumonia, epilepsy, fever, convulsion, diarrhea, headache and influenza.	[27 - 34]
<i>Ocimum Kilimandscharicum</i>	India, Thailand, Ethiopia, Tanzania, Kenya, Uganda, Sudan, Ethiopia	Leaves	It is employed in the treatment of cough, cold, measles, abdominal pain, measles, diarrhea and diarrhea. It is used as an insect repellent and for storage pest control.	[35 - 37]
<i>Ocimum sanctum</i>	Asia, Africa, Malaysia, Australia, United Arab Emirates	Leaves	It is used traditionally for the management and treatment of several ailments such as headache, fever, cough, common cold, flu, sore throat, colic pain, asthma, diarrhea, digestive disorders, bronchitis, influenza, insomnia, hepatic diseases, arthritis, and malaria fever. It is also used as an antidote for scorpion sting and snake bite	[38 - 40]

Table 1. The tradomedic al uses of different *Ocimum* species

Phytochemical Studies

Ethnomedicinal plants contain complex plant chemicals also known as phytochemicals. These phytochemicals are made up of several compounds. The enormous information revealed by the ethnomedicinal applications of *Ocimum species* spurred scientific investigation into its chemical constituents. These studies have led to the identification and isolation of divers phytochemicals from different plant parts of *Ocimum* species. These include alkaloids, terpenoids, organic acids, tannins, flavonoids, coumarins, quinones, polyphenols, saponins, and their derivatives.

Ocimum americanum

Phytochemical analysis carried-out by Dhale *et al*^[41] showed that *Ocimum americanum* is rich in alkaloids, phenolic compounds, tannins, lignin, saponins, flavonoids, terpenoid and anthraquinone. Alciléia *et al*^[42] also gave a similar report, and as well documented compounds such as α -pinene, camphene, sabinene, β -pinene, β -mircene and other compounds which have been identified and isolated from its essential oil.

Ocimum basilicum

Sanni *et al*^[43] and Ismail^[44] reported the presence of carbohydrate, tannins, cardiac glycosides, flavanoids, terpenes and steroids in *Ocimum basilicum* leaves. Using a GC-MS, Andrew *et al*^[45] carried-out a chemical analysis which showed the presence of α -Pinene, β -Myrcene, 4-Hexen-1-ol, acetate and other important bioactive compounds in its leaf essential oil. Sarfaraz *et al*^[46] also documented the presence of these compounds in a report on their analysis.

Ocimum gratissimum

A phytochemical report documented by Offiah and Chikwendu^[47] revealed the presence of tannins, steroids, triterpinoids and carbohydrates in *ocimum gratissimum*. Prabhu *et al*^[48] also showed the presence of alkaloids, tannins, flavonoids and oligosaccharides. Pandey and Chowdhury,^[49] Matasyoh *et al*,^[50] Kéita *et al*,^[51] and Jirovetz *et al*^[52] have also documented specific compounds which have been isolated from *Ocimum gratissimum*, this have been summarized in table 2 and some of the structures shown in figure 1.

Ocimum Kilimandscharicum

According to a report by Tewari *et al*,^[53] alkaloids, glycosides, saponins, flavanoids, phenols, carbohydrates, steroids, protein and amino acids are among the phytochemicals present in *Ocimum Kilimandscharicum*. Charles and Simon^[54] in their work isolated seventeen compounds from its essential oil and this includes α - pinene, Camphene, β -pinene and Eugenol [table 2].

Ocimum sanctum




In a research work published by Xia *et al*,^[55] carbohydrates, terpenoids and alkaloids were present in the ethanolic extract of *Ocimum sanctum*. Similarly, Devendran and Balasubramanian^[56] also reported the presence of flavanoids, cardiac glycosides, steroids, saponin and tannins, as well as carbohydrates, terpenoids and alkaloids. Furthermore, using a GC-MS, analysis of the hydro-alcoholic extract showed the presence of several compounds including Eugenol and Caryophyllene.

Reported compounds isolated from the various *Ocimum* species have been outlined in table 2, while some of the structures are shown in figure 1.

Species	Compounds Isolated	Reference
<i>Ocimum americanum</i>	α -pinene, Camphene, Sabinene, β -pinene, β -myrcene, Limonene, 1,8 cineole, linalool oxide, Fenchone, Linalool, Camphor, Borneol, 4-terpineol, α -Terpineol, cis-piperitol, Estragole, Eugenol, β -selinene, β -caryophyllene, bicyclogermacrene, β -bisabolene.	[42]
<i>Ocimum basilicum</i>	α -Pinene; β -Myrcene; 4-Hexen-1-ol, acetate; 4 Eucalyptol; cis-Linalooloxide; 1,6-Octadien-3-ol, 3,7-dimethyl; Methyl ethyl cyclopentene; L-Menthone; L-(β)-menthol; Estragole; N-Cyano-3-methylbut-2-enamine; Citral; Cyclohexene, 4-methyl-1-(1-methyl-ethyl); Phenol, 2,3,5-trimethyl; Eugenol; Formic acid, cyclohexyl ester; Copaene; cis-7,10,13,16-Docosatetraenoic acid, methyl ester; Neoisolongifolene; trans- α -Bergamotene; Alloaromadendrene; Humulene, beta.-copaene; beta.-Bisabolene; cis-muurola-3,5-diene;	[45 - 46]

	<p>cis-.alpha.-Bisabolene; Nerolidol; trans-4-Methoxycinnamaldehyde; Benzeneacetic acid,.alpha.-hydrox; Phenylethanolamine; 3-Methyl-2-phenylindole; N-Benzyl-N-ethyl-p-isopropylbenzamide</p>	
<i>Ocimum gratissimum</i>	<p>Eugenol, methyl eugenol, cis-ocimene, trans-ocimene, pinene, camphor, germacrene-D, trans-caryophyllene, farnesene and l-bisabolene, Thymol, methyl chavical, linalool, limonene, methyl eugenol, β-caryophyllene, farnesene, α-terpineol, methyl isoeugeneol, geraniol, α-copaene, bisabolol, fenchone, cubenene, camphene, T-cadanol, sabinene, myrcene, β-bisoboline, α-humelene and β-elemene.</p>	[47 -52]
<i>Ocimum Kilimandscharicum</i>	<p>β-pinene, Myrcene, 1,8-cineole, Limonene , Terpinen-4-ol, α-terpineol, Bornyl acetate, α-pinene, Camphene, Eugenol, β-caryophyllene, α-humu lene, Linalool, Camphor, γ-muurolene, Germacerene B, Epi-α-cadinol</p>	[53-54]
<i>Ocimum sanctum</i>	<p>Eugenol; Cyclohexane, 1,2,4-triethyl- ; Caryophyllene; 10-Heptadecen-8-ynoic acid; Cyclopentane, cyclopropylidene-; Z,Z-4,16-Octadecadien-1-ol acetate; Benzene methanamine, N,N-α,4-tetramethyl-; 3',8,8'-trimethoxy-3-piperidyl-2,2'-binaphthalene1,1',4,4'-tetron e; Octadecane, 1,1-dimethoxy-; Pentanedinitrile, 2-methyl-</p>	[55-56]

Table 2. Compounds isolated from different *Ocimum* species












Camphor

Eugenol

 β -Selinene

Estragol

Camphene

Trans β -Ocimene**Figure 1.** Structures of some compounds identified/isolated from *Ocimum* species

Pharmacology

Several pharmacological studies have confirmed/establish the efficacy of *Ocimum* species for their various therapeutic applications.

Ocimum americanum

Ocimum americanum have been documented to elicit significant antifungal activity,^[57 - 59] gastric cyto-protective and antiulcer effect against acute gastric ulcer causative organism,^[57 - 60] wound healing activity,^[57] larvicidal and insecticidal activity.^[61] The essential oils also have effects such as cytotoxic, antimicrobial and mosquito repellent effect.^[59, 62] The antioxidant activity of *Ocimum americanum* has also been documented.^[63 - 64] Sutili et al reported that *Ocimum americanum* whole plant have a significant hemolytic effect.^[65] Sethi et al documented the antifungal activity of the plant,^[66] while Priscila et al reported the bactericidal activity.^[67] The anti-microbial activity against gram positive and gram negative bacteria have been reported by Thawebon et al.^[68] Aluko et al also reported its anti-oxidant effect.^[69] Sripriya et al reported the significant analgesic and anti-inflammatory activities of the leave in laboratory mice.^[70] According to Sinitha et al, *Ocimum americanum* elicits immune-modulatory effect,^[71] while Lenise et al have also documented its anesthetic activity.^[72] Aluko et al carried-out a study to evaluate the hepatoprotective activity of *Ocimum americanum* leaves in laboratory rats with paracetamol - induced liver damage, and reported it to be having an hepatoprotective effect.^[73]

Ocimum basilicum

According to reports by Politeo et al and Ramesh et al, *Ocimum basilicum* is rich in antioxidants compounds and hence have a high antioxidant activity.^[74 - 75] Dwivedi et al stated that the Solvent extracts of the plant elicits significant activity at cellular level including inhibition of HIV-1 reverse transcriptase and anti-aggregatory effect on blood platelets.^[76] Clinical studies reported by Siurin et al and Niture et al revealed that *Ocimum basilicum* volatile oil have characteristic antioxidant activities.^[77 - 78] Bravo et al also reported the protective activity of the ethanolic extract of *Ocimum basilicum* against mutagenesis and oxidative DNA damage, with accompanying decrease in cholesterol synthesis and accumulation of lipid in human macro-phages.^[79] Report by Wannissorn et al and Ahmet et al reveals that *Ocimum basilicum* exhibits antibacterial effect against a several bacteria including *C. perfringens*, *Salmonella* spp., *C. jejunii* and *E. coli*.^[80 - 81] Also, Opalchenova and Obreshkova revealed that the plant aerial parts volatile oil elicits significant activity against drug immune clinical isolates of *Pseudomonas*, *Enterococcus* and *Staphylococcus*.^[82] Patel et al have documented the significant antimicrobial effects of *Ocimum basilicum* from their study.^[83] In vitro studies by Yamasaki et al revealed that *Ocimum basilicum* exhibits marked inhibitory actions against cytopathogenicity induced by HIV-1 in MT-4 cells.^[84] Renzulli et al reported that rosmarinic acid (a compound present in *Ocimum basilicum*) has a significant inhibitory activity against inflammatory processes. In vitro studies reveal that rosmarinic reduces the production of oxygen species, as well as inhibits protein and DNA synthesis induced by mycotoxins, hence prevents cell death.^[85] In vitro study on P388 and KB cell lines by Manosroi et al revealed that the essential oil *Ocimum basilicum* elicited moderate anti-proliferative activity compared to the standard drug, 5-fluorouracil.^[86] Mohan et al reported the hypolipidemic and hypoglycemic activity of aqueous extract of *Ocimum basilicum* in laboratory rats. They observed marked reduction in serum lipid, lipid peroxidation products, blood glucose level, as well as improvement in glucose tolerance.^[87]

Ocimum gratissimum

The antibacterial and antifungal efficacy of *Ocimum gratissimum* has been reported in various studies.^[88 - 89] Pharmacological study of the aqueous extract revealed it has analgesic effect in mice, stomach strip blocking effect in rat and inhibitory effect on rabbit jejunum pendular movement.^[90]

Report by Iwalokun et al revealed that the oil elicits a significant inhibitory activity against the virulent strains of *Shigella*, *E. coli* and *Salmonella*.^[91] The hexane fraction exhibits a highly significant antimicrobial activity against *K. pneumonia* and *V. cholera*.^[92] It is significantly active against *P. aeruginosa*, *S. typhi*, *V. cholera* and *N. gonorrhoea*.^[93] It has a good synergic activity with ampicillin, eliciting highly significant antibacterial effects against *E. coli* and *P. mirabilis*, and also possess similar synergic activity with nystatin and ketoconazole, eliciting significant antifungal effect against *C. albicans*.^[94] The anti-diarrheal activity of the leaf extract was established in a study by Ilori et al which used the disc diffusion and tube dilution methods to ascertain its effect against bacteria *S. typhi*, *E. coli*, *S. dysenteriae*, *P. shigelloides* and *A. sobria*, and it proved to be a significant antibacterial agent.^[95] Report by Pratheeba et al also revealed that the chloroform leaf extract had a good insecticidal effect against filariasis causing mosquito vector *Culex quinquefasciatus*.^[96] The plant leave have been shown to possess a broad-spectrum anti-bacterial activity against *P. mirabilis*, *E. coli* and *S. aureus*.^[97] The methanolic leaf extract have been reported to be active against nicotine toxicity due to its ability to decrease free radical generation and lipid-protein damage, and its antioxidant activity.^[98] The aqueous leaf extract have been reported to show significant hypoglycaemic effect, reducing lipid MDA, LDL- cholesterol and triacylglycerol levels^[99] and also, caused a marked decrease in the plasma glucose level in laboratory rats in streptozotocin induced diabetic model.^[100] The aqueous leaf extract have been observed to inhibit proliferation, chemotaxis, induction of COX-2 protein and morphogenesis and as well decrease size of breast cancer tumor.^[101] The dichloromethane leaf extract in an in vitro study inhibited myeloid leukemia,^[102] hence, may be useful in treating human cancer. The methanolic crude leaf extract have been documented to be having haematinic and haemopoietic activity in laboratory rats.^[103] Increase in erythrocytes, hemoglobin packed cell volume, thrombocytes and neutrophils count, as well as increase in platelet distribution width, mean platelet volume, and platelet-large cell ratio (P-LCR) have been observed to be elicited by the leaf extract.^[104] Reduction in the serum total protein and urea level, packed cell volume, neutrophils and hemoglobin, and increase uric acid, in total acid, prostatic acid, phosphatases, leucocytes and lymphocytes have also been reported to be elicited by the plant.^[105] Its leaf essential oil have been reported to show both fungistatic and fungicidal activity at different concentration against *A. alternata*, *S. rolfsii* and *C. capsici*.^[106]

Ocimum kilimandscharicum

Hakkim et al evaluated the antioxidant potential of the Methanolic leaf extract of *Ocimum kilimandscharicum* using an in vitro model. They reported it to exhibit a moderate activity compared to butylated hydroxyl anisole (BHA).^[107] Mwangi et al also carried-out an experiment to examine the antinociceptive activity of the alcoholic leaf extract of *Ocimum kilimandscharicum* in mice using the tailflick model. The outcome revealed it to be a significant antinociceptive agent.^[108] Based on their in-vitro antiplasmodial study using dichloromethane extract of leaves and twigs of *Ocimum kilimandscharicum*, Owuor et al reported significant activity against both chloroquine resistant and chloroquine sensitive *Plasmodium* parasites,^[109] while Runyoro et al reported that the oil elicited larvicidal activity on *Culex quinquefasciatus* larvae.^[110] Using castor-oil induced diarrhoea model and castor oil induced enteropooling assay in rats; and charcoal meal test/intestinal motility test in mice, Serin et al demonstrated that the aqueous leaf extract has a significant anti-diarrheal activity.^[111] Kumar et al reported the significant antimicrobial activity of volatile oil of whole plant of *Ocimum kilimandscharicum* against Gram negative bacteria (*E. coli*, *E. cloacae*, *Vibrio cholera*, *S. dysenteriae*, *P. aeruginosa*, *Klebsiella sp*),^[112] while Shinde et al also reported its effectiveness against Gram positive bacteria (*S. aureus*, *S. epidermidis*, *S. mutans*, *S. viridians*, *B. subtilis*, *M. luteus*).^[113] Monga et al also reported the antimelanoma and radioprotective effect of alcoholic aqueous extract in mice.^[114] Jambere et al carried-out a laboratory test on the leaves of *Ocimum kilimandscharicum* against the following insects *Rhyzopertha dominica*, *Sitophilus zeamais*, *Sitotroga cerealella* in maize and sorghum grains to evaluate its insecticidal activity. The reported experimental outcome showed it to be an effective insecticidal agent.^[115] Mahesh et al has also documented the significant wound healing activity of the aqueous extract as a result of the experiment they carried out on laboratory rats, using the

excision, incision and dead space wound model.^[116]

Ocimum sanctum

Ocimum sanctum has been chronicled and described to be having several pharmacological activities. Report by Rahman et al reveals that *Ocimum sanctum* shows antimicrobial activity against wide range of bacteria including *Staphylococci* sp., *E. coli*, *Shigella* sp., *S. aureus*, *Enterobacteria* sp., *P. aeruginosa*, *S. typhi*, *Staphylococci* sp., *C. albicans*, *K. pneumonia*, *M. tuberculosis* and *M. pyogenes*.^[117] According to an in vitro study by Farivar et al, *Ocimum sanctum* extract proved to be an effective anti-tuberculosis agent.^[118] Khan et al reported that aqueous, hexane, chloroform, n-butanol extracts and essential oil of *Ocimum sanctum* showed significant antifungal activity against several fungi including *A. solani*, *C. guillermondii*, *C. capsici*, *Curvularia* sp., *F. solani*, *H. oryzae* and *A. flavus*.^[119-120] Kelm et al also reported that the methanolic, hydroalcoholic and aqueous extracts of *Ocimum sanctum* elicits significant antioxidant activity, both *in vitro* and *in vivo*.^[121] According to Geetha et al, Oral intake of *Ocimum sanctum* gives significant protection of liver and aortic tissue from hypercholesterolemia induced peroxidative damage.^[122] Siva et al documented that the Oral administration of *Ocimum sanctum* extract significantly reduced blood sugar level in streptozotocin-induced and glucose-fed hyperglycemic diabetic rats. A different study by Gholap and Kar also showed that *Ocimum sanctum* reduced the serum concentration of cortisol and that of glucose and as well elicited antiperoxidative activity.^[123] Aruna et al in their study revealed that *Ocimum sanctum* leaves significantly reduced the squamous cell hematoma incidences and carcinoma in experimental rats.^[124] Result from a study by Ganasoundari et al revealed that *Ocimum sanctum* aqueous leaf extract inhibited hydroxyl (OH) radical-induced deoxyribose degeneration significantly. They also showed the synergic activity WR-2721 and *Ocimum sanctum* which produced a better effect against OH radical activity when compared with the individual agents alone.^[125] Shetty et al performed an experiment to evaluate the activity of *Ocimum sanctum* aqueous leaf extract on tumor necrosis factor-Alfa (TNF-Alfa) in laboratory rats, using the excision model. The result showed that *Ocimum sanctum* extract increased the rate of epithelization and wound contraction, hence possesses a significant wound healing effect.^[126] In antiulcer study carried-out by Govind et al in rats using aspirin, indomethacin, alcohol (ethanol 50%), histamine, reserpine, serotonin or stress-induced ulcers models, the oil of *Ocimum sanctum* showed significant antiulcer activity.^[127] Mediratta et al, reported that steam distilled leave extract of *Ocimum sanctum* caused modification in laboratory rat humoral immune response and this could be attributed to antibody production, release of mediators of hypersensitivity reactions and tissues responses to mediators in the target organs.^[128] Godhwani et al in a separate experiment also demonstrated the immunomodulatory effect of *Ocimum sanctum* using widal and sheep erythrocyte agglutination tests.^[129] Sridevi et al showed that *Ocimum sanctum* has significant benefits when employed in the treatment of asthma and other related conditions. They also reported that *Ocimum sanctum* has the potential to cause mast cell stabilization, suppression of IgE and inhibition of release of inflammatory mediators, hence, may be the cause of these effects.^[130] According to Ravindran et al, *Ocimum sanctum* normalizes noise stress induced alteration in neurotransmitter levels due to, and this is a proof of its anti-stress activity.^[131] Asha et al showed the significant antihelmentic activity of *Ocimum sanctum* essential oil using the caenorhabditis elegans model.^[132] Different extracts of stem and leaves of *Ocimum sanctum* were subjected to experimental studies for anticonvulsant activity by maximal electroshock model using phenytoin as standard, and was observed that ethanol and chloroform extract of leaf and stem produced significant preventive effect against toxic convulsions induced by trans corneal electroshock.^[133] *Ocimum sanctum* have also been shown to have strong cardio protective effect against myocardial agents. An experiment by Sood et al revealed that *Ocimum sanctum* elicited significant protection against isoproterenol-induced myocardial necrosis in laboratory rats by increasing the activity endogenous antioxidants.^[134] Joshi et al reported that *Ocimum sanctum* L. alcoholic extract ameliorated scopolamine-induced amnesic effect as well as memory deficits caused by aging in laboratory mice. *Ocimum sanctum* both increased significantly step-down latency (SDL) and acetylcholinesterase inhibition. This may be useful in the treatment of dementia, alzheimer's disease and other cognitive disorders.^[135] *Ocimum sanctum* have been

documented to have significant antidote effect against dog bite, snake bite, scorpion bite and insect bites.^[136]

Conclusion

Ocimum species have been employed traditional for therapy globally. The species reviewed are the most widely used from the *Ocimum* genus in traditional medicine and they are also the most researched for intended use in conventional therapeutic practice. These species contain several compounds which are responsible for the numerous pharmacological activities elicited by them including antimalarial, anticancer, antidiabetic, antiinflammatory and antioxidant effects.

References

1. Ashraf K, Haque MR, Amir M, Ahmad N, Ahmad W, Sultan S, Ali Shah SA, Mahmoud AA, Mujeeb M, Bin Shafie MF. An overview of phytochemical and biological activities: *Ficus deltoidea* Jack and other *Ficus spp.* J Pharm Bioall Sci 2021;13:11-25.
2. Bunawan H, Amin NM, Bunawan SN, Mohd Noor N. *Ficus deltoidea* jack: a review on its phytochemistry and pharmacological importance. Evi-Based Comp Alter Med 2014;2:1-8.
3. Chinedu E, Arome D, Ameh SF, Ameh GE. Evaluation of the anti-proliferative and cytostatic effect of *Citrus sinensis* (orange) fruit juice. Int J App Basic Med Res 2014;4:20-2.
4. Chowdhury T, Mandal A, Roy SC, De Sarker D. Diversity of the genus *Ocimum* (Lamiaceae) through morpho-molecular (RAPD) and chemical (GC-MS) analysis. J Gen Eng Biotech 2017;15:275-286.
5. Hereman S. Paxton's Botanical Dictionary. Bradbury Evans and Co., London, 1868.
6. Simpson BB, Corner OM. Economic Botany-Plants in our World, McGraw-Hill Book Company, Hamburg, 1986.
7. Pushpangadan P, Bradu BL. In: Chadha KL, Gupta R (Eds.), Advances in Horticulture, Malhotra Publishing House, New Delhi, 1995.
8. Paton A, Harley RM, Harley MM. In: Hiltunen R, Holm Y. (Eds.), *Ocimum: An Overview of Classification and Relationship*, Harwood Academic Publishers, Amsterdam, 1999.
9. Sandeep P. The phytochemical constituents, pharmacological and traditional uses of *Ocimum gratissimum* L in tropics. Indo Am P Sci 2017;4(11):4234 - 4242.
10. Nwanjo HU, Oze GO. Hypolipidaemic and antioxidant properties of *Ocimum gratissimum* on diabetic rats. Plant Prod Res J 2007;11:1-4.
11. Rao SA, Vijay Y, Deepthi T, Lakshmi CS, Rani V. Antidiabetic effect of ethanolic extract of leaves of *Ocimum sanctum* in alloxan induced diabetes in rats. Int J Basic Clin Pharmacol 2013;2:613-616.
12. Maddi R, Amani P, Bhavitha S, Gayathri T, Lohitha T. A review on *Ocimum* species: *Ocimum americanum* L., *Ocimum basilicum* L., *Ocimum gratissimum* L. and *Ocimum tenuiflorum* L. Int J Res Ayurveda Pharm 2019;10(3):41-48.
13. Nyarko AK, Asare-Anane H, Ofosuhene M, Addy ME. Extract of *Ocimum canum* lowers blood glucose and facilitates insulin release by isolated pancreatic beta-islet cells. Phytomed 2002;9:346-351.
14. Obeng-Ofori D, Reichmuth CH, Bekele AJ, Has-sanali A. Toxicity and protectant potential of camphor, a major component of essential oil of *Ocimum kilimandscharicum*, against four stored product beetles. Int J Pest Manag 1998;44:203-209.
15. Pattnaik S, Chand PK. In vitro propagation of the medicinal herbs *Ocimum americanum* L. syn. O. canum Sims.(hoary basil) and *Ocimum sanctum* L.(holy basil). Plant Cell Reports 1996; 15(11):846-850.
16. Watt JM, Gerdinat M, Breyer-Brandwijk K. Medicinal and poisonous plants of southern and eastern africa,. Second Edition. E & S Lwinstone Ltd; 1962.
17. Githinji CW, Kokwaro JO. Ethnomedicinal study of major species in the family Labiatae from Kenya. Journal of Ethnopharmacology 1993; 39(3):197-203.
18. Kokwaro J. Medicinal Plants of East Africa. Second Edition. Kenya Literature Bureau;

1993.

19. Vidhya E, Vijayakumar S, Rajalakshmi S, Kalaiselvi S, Pandiyan P. Antimicrobial activity and phytochemical screening of *Ocimum americanum*L extracts against pathogenic microorganisms. *Acta Ecologica Sinica* 2020; 40(3):214-220.
20. Harnafia H, Azizb M, Amrania S. Sweet basil (*Ocimum basilicum* L.) improves lipid metabolism in hypercholesterolemic rats. *Euro e-Journal Clin Nutri Metabol.* 2009; 4:181-6.
21. Umar A, Zhou W, Abdusalam E, Tursun A, Reyim N, Tohti I, Moore N. Effect of *Ocimum basilicum* L. on cyclo-oxygenase isoforms and prostaglandins involved in thrombosis. *J Ethnopharmacol* 2014;152: 151-5.
22. Tsai KD, Lin BR, Perng DS, Wei JC, Yu YW, Cherng JM. Immunodulatory effects of aqueous extract of *Ocimum basilicum* (Linn.) and some of its constituents on human immune cells. *J Med Plants Resear* 2011;5:1873-83.
23. Opalchenova G, Obreshkova D. Comparative studies on the activity of basil-an essential oil from *Ocimum basilicum* L. against multidrug resistant clinical isolates of the genera *Staphylococcus* and *Pseudomonas* by using different test methods. *J Microbiol Methods.* 2003; 54: 105-10.
24. Caceres A, Cano O, Samayoa B, Aguilar L. Plants used in Guatemala for the treatment of gastrointestinal disorders. 1. Screening of 84 plants against enterobacteria. *J Ethnopharmacol* 1990;30:55-73.
25. Ngassoum MB, Ousmaila H, Ngamo LT, Maponmetsem PM, Jirovetz L, Buchbauer G. Aroma compounds of essential oils of two varieties of the spice plant *Ocimum canum* Sims. From northern Cameroon *J Food Comp Anal* 2004;17:197-204.
26. Bassole IHN, Guelbeogo WM, Nebie R, Costantini C, Sagnon NF, Kabore ZI, Traore SA. Ovicidal and larvicidal activity against *Aedes aegypti* and *Anopheles gambiae* complex mosquitoes of essential oils extracted from three spontaneous plants of Burkina Faso. *Parasitol* 2003;45:23-26.
27. Adebolu TT, Oladimeji SA. Antimicrobial activity of leaf extracts of *Ocimum gratissimum* on selected diarrhoea causing bacteria in Southwestern Nigeria. *Afri J Biotechnol* 2005;4:682-684.
28. Aguiyi JC, Obi E, Gang SS, Igweh A. Hypoglycaemic activity of *Ocimum gratissimum* in rats. *Fitoterapia* 2000;71: 444-446.
29. Akinyemi O, Mendie U, Smith S, Oyefolu A, Coker A. Screening of some medical plants for anti-salmonella activity. *J Herb Pharmacother* 2004;5:45-60.
30. Gbolade AA. Inventory of antidiabetic plants in selected districts of Lagos State, Nigeria. *J Ethnopharmacol* 2009;121:135-139.
31. Lopez P, Sanchez C, Batlle R, Nerin C. Solid and vapor-phase antimicrobial activities of six essential oils: Susceptibility of selected food-borne bacterial and fungi strains. *J Agri Food Chem* 2005;53:6939-6946.
32. Odukoya A, Ilori O, Sofidiya M, Aniunoh O, Lawal B, Tade I. Antioxidantactivity of Nigerian dietary spices. *EJEAF Chem* 2005;46:1086-1093.
33. Ezekwesili CN, Obiora KA, Ugwu OP. Evaluation of anti-diarrhoeal property of crude aqueous extract of *Ocimum gratissimum* L.(Labiatae)in rats. *Biokemistri* 2004;16:122-131.
34. Freire MM, Marques OM, Costa M. Effects of seasonal variation on the central nervous system activity of *Ocimum gratissimum* L. essential oil. *J Ethnopharmacol.* 2006; 105:161-6.
35. Ofori DO, Reichmuth CH, Bekele AJ, Hassanali A. toxicity and protectant potential of camphor, a major component of essential oil of *ocimum kilimandscharicum*, against four-stored product beetles. *Int J Pest Manag* 1998;44(4):203-209.
36. Bekele J, Hasannali A. Blend effects in the toxicity of the essential oil constituents of *ocimum kilimandscharicum* and *Ocimum kenyense* (Labiatae) on two poat-harvest pests. *Phytochem* 2001;57:385-391.
37. Misra RC, Das G. *Ocimum kilimandscharicum* Guerke (Lamiaceae): A new distributional record for peninsular India with focus on its economic potential. *Proc Natl Acad Sci India Sect B Biol Sci* 2016;86:795-803.
38. Pandey G, Madhuri S. Pharmacologocal Activities of *Ocimum sanctum* (Tulsi): A Review. *Int. J. Pharm. Sci. Rev. Res* 2010;5(1):61-66.

39. Ogen S, Mas'ud H, Setiawan K, Muchammad Y, Ngakan MRW. Anthelmintic Activity of Basil Leaves (*Ocimum sanctum* Linn.) Infusion Against *Ascaris suum* In Vitro. *J Parasite Sci* 2017;1(2):47-50.

40. Rai V, Mani UV, Iyer UM. Effect of *Ocimum sanctum* leaf powder on blood lipoproteins, glycated proteins and total Amino acids in patients with Non-insulin-dependent Diabetes Mellitus. *J Nutri Environ Med* 2009;7: 113-8.

41. Dhale DA, Birari AR, Dhulgande GS. Preliminary Screening of Antibacterial and Phytochemical Studies of *Ocimum americanum* Linn. *J Ecobiotechnol* 2010;2(8):11-13.

42. Alciléia NY, Renata G, Aureo TY, Expedito LS, Saulo ES, Marcio JD, Márcia MOD, Ciomar AB, Roberto KNC. Anti-inflammatory Activity of *Ocimum americanum*L. Essential Oil in Experimental Model of Zymosan-Induced Arthritis. *Am J Chin Med* 2013; 41(4): 913-926

43. Sanni S, Onyeyili PA, Sanni FS. Phytochemical Analysis, Elemental Determination and Some *in vitro* Antibacterial Activity of *Ocimum basilicum* L. Leaf Extracts. *Resear J Phytochem*, 2008;2:77-83.

44. Ismail M. Central properties and chemical composition of *Ocimum basilicum* essential oil. *Pharmaceut Bio* 2006; 44(8):619-626.

45. Andrew BF, Felicitas EM, Emrobawansan MI, Anthony JA, Voster M. Phytochemical Constituents and Antioxidant Activity of Sweet Basil (*Ocimum basilicum* L.) Essential Oil on Ground Beef from Boran and Nguni Cattle. *Int J Food Sci* 2019;2019:1- 8.

46. Sarfaraz KM , Fazal UR , Muhammad SK, Said G, Naveed A, Ghulam M, Khalid U. Phytochemical Constituents and Pharmacological Activities of Sweet Basil-*Ocimum basilicum* L. (Lamiaceae). *Asian J Chem* 2011;23(9):3773-3782

47. Offiah VN, Chikwendu UA. Antidiarrhoeal effects of *Ocimum gratissimum* leaf extract in experimental animals. *J Ethnopharmacol* 1999;68:327-30.

48. Prabhu KS, Lobo R, Shirwaikar AA, Shirwaikar A. *Ocimum gratissimum*: A Review of its Chemical, Pharmacological and Ethnomedicinal Properties. *The Open Complem Med J* 2009;1:1-15.

49. Pandey AK, Chowdhury AR. Composition of the essential oil of *Ocimum gratissimum* grown in Madhya Pradesh. *J Med Aromat Plant Sci* 2000; 22-23, 26-8.

50. Matasyoh LG, Josphat CM, Francis NW, Miriam GK, Anne WTM, Titus KM. Chemical composition and antimicrobial activity of the essential oil of *Ocimum gratissimum* L. growing in Eastern Kenya. *Afr J Biotech* 2007; 6: 760-5.

51. Kéita SK, Vincent C, Jean-Pierre S, Bélanger A. Essential oil composition of *Ocimum basilicum* L., *O. gratissimum* L. and *O. suave* L. in the Republic of Guinea. *Flav Fragr J* 2000; 5:339-41.

52. Jirovetz L, Buchbauer G, Ngassoum MB, Ngamo LT, Adjoudji O. Combined investigation of the chemical composition of essential oils of *Ocimum gratissimum* and *Xylopia aethiopica* from Cameroon and their insecticidal activities against stored maize pest *Sitophilus zeamais*. *Ernähr* 2005; 29:55-60.

53. Tewari D, Pandey HK, Sah AN, Meena HS, Manchanda A. Pharmacognostical and biochemical investigation of *Ocimum kilimandscharicum* plants available in western Himalayan region. *Asian Journal of Plant Science and Research*, 2012, 2 (4):446-451

54. Charles DJ, Simon JE, Essential oil constituents of *Ocimum kilimandscharicum* Guerke, *J Essential Oil Resear* 1992;4:125-128.

55. Xia KZ, Perveen N, Khan NH. Phytochemical analysis, antibacterial and antioxidant activity determination of *Ocimum sanctum*. *Pharm Pharmacol Int J* 2018;6(6):490-497.

56. Devendran G, Balasubramanian U. Qualitative phytochemical screening and GC-MS analysis of *Ocimum sanctum* L. leaves. *Asian J Plant Sci Resear*, 2011;1(4):44-48

57. Sutili FJ, Velasquez A, Pinheiro CG, Heinzmann BM, Gatlin III DM, Baldisserotto B. Evaluation of *Ocimum americanum* essential oil as an additive in red drum (*Sciaenops ocellatus*) diets. *Fish and Shell fish Immunol* 2016; 56:155-161.

58. Vidhya E, Vijayakumar S, Rajalakshmi S, Kalaiselvi S, Pandiyan P. Antimicrobial activity and phytochemical screening of *Ocimum americanum*L extracts against pathogenic microorganisms. *Acta Ecologica Sinica* 2020; 40(3):214-220.

59. Zengin G, Ferrante C, Gnapi DE, et al. Comprehensive approaches on the chemical

constituents and pharmacological properties of flowers and leaves of American basil (*Ocimum americanum*L). Food Resear Int 2019; 125:108610.

60. Carović-Stanko K, Orlić S, Politeo O, et al. Composition and antibacterial activities of essential oils of seven *Ocimum* taxa. Food Chem 2010; 119(1):196-201.

61. Shadia E, El-Aziz A, Omer EA, Sabra AS. Chemical composition of *Ocimum americanum* essential oil and its biological effects against *Agrotis ipsilon* (Lepidoptera: Noctuidae). Research Journal of Agri Biol Sci 2007;3(6):740-747

62. Runyoro D, Ngassapa O, Vagionas K, Aligiannis N, Graikou K, Chinou I. Chemical composition and antimicrobial activity of the essential oils of four *Ocimum* species growing in Tanzania. Food Chem 2010; 119(1):311-316.

63. Chokechajaroenporn O, Bunyapraphatsara N, Kongchuensin S. Mosquito repellent activities of *Ocimum* volatile oils. Phytomed 1994;1(2):135-139.

64. Madhiyazhagan P, Murugan K, Kumar AN, Nataraj T. Extraction of mosquitocidals from *Ocimum canum* leaves for the control of dengue and malarial vectors. Asian Pac J Trop Dis 2014; 4:S549-S555.

65. Sutilifj, Velasquez, Pinheiro CG, Baldesserotto, Evaluation of *Ocimum americanum* essential oil as an additive in red drum (*Sciaenops ocellatus*) diets, Fish shellfish imunol 2016 Sep; 56:155-161.

66. Sethi S, Prakash Om, Chandra M, Punetha H, Pant A K. Antifungal activity of essential oils of some *Ocimum* species collected from different locations of Uttarakhand, Indian Journal of Natural Product Resources 2013; 4(4):392-397.

67. Priscila RN, Vieira Selene Mde, Morais Francisco HQ, Bezerra Pablito Augusto Travassos Ferreira Írvila, ROliveira Maria Goretti V. Silva. Chemical composition and antifungal activity of essential oils from *Ocimum* species. Industrial Crops and Products 2014 April; 55: 267-271

68. Thaweeboon S, Thaweeboon B, In vitro antimicrobial activity of *Ocimum americanum*L. essential oil against oral microorganisms. The Southeast Asian Journal of Tropical Medicine and Pub Heal 2009; 40(5):1025-1033.

69. Aluko BT, Oloyede OI, Afolayan AJ, Polyphenolic contents and free radical scavenging potential of extracts from leaves of *Ocimum americanum*L. Pakistan Journal of Biological Sciences 2013; 16(1):22-30.

70. Sripriya D, Venkanna L, Estari Mamidala. Analgesic and anti-inflammatory effects of *Ocimum americanum*(Linn) In Laboratory Animals. IJPSR 201; 2(8) 2121-2125.

71. Sunitha, Nasreen begum. Immunomodulatory activity of methanolic extract of *Ocimum americanum*seeds. Journal of Pharmaceutical and Biomedical Analysis 2011; 2(1):33-38.

72. Lenise de Lima Silva, Quelen Iane Garlet, Gessi Koakoski, Murilo Sander de Abreu, Carlos Augusto Mallmann , Bernardo Baldisserotto , Leonardo José Gil Barcellos and Berta Maria Heinzmann. Anesthetic activity of the essential oil of *Ocimum americanum*in Rhamdia quelen (Quoy & Gaimard, 1824) and its effects on stress parameters. Neotrop Ichthyol 2015; 13(4):715-722.

73. Aluko BT, Oloyede OI, Afolayan AJ. Hepatoprotective activity of *Ocimum americanum*L leaves against paracetamol - induced liver damage in rats. American Journal of Life Sciences 2013 April; 1(2):37-42.

74. Politeo O, Jukic M, Milos M. Chemical composition and antioxidant capacity of free volatile aglycones from basil (*Ocimum basilicum* L.) compared with its essential oil. Food Chem. 2007; 101, 379-85.

75. Ramesh B, Satakopan VN. 'In vitro' Antioxidant activities of *Ocimum* species; *Ocimum basilicum* L. and *Ocimum sanctum*. J. Cell and Tissue Res. 2010; 10: 2145-50.

76. Dwivedi C, Abu-Ghazaleh A. Chemo-preventive effects of sandalwood oil on skin papillomas in mice. Eur. J. Cancer Prev. 1997; 6: 399-401.

77. Siurin SA. Effects of essential oil on lipid peroxidation and lipid metabolism in patients with chronic bronchitis. Klin Med (Mosk). 1997; 75: 43-5.

78. Niture SK, Rao US, Srivenugopal KS. Chemo preventative strategies targeting the MGMT repair protein; augmented expression in human lymphocytes and tumor cells by ethanolic and aqueous extracts of several Indian medicinal plants. Int. J. Oncol. 2006; 29: 1269-78.

79. Bravo E, Amrani S, Aziz M, Harnafi H, Napolitano M. *Ocimum basilicum* methanolic extract decreases cholesterol synthesis and lipid accumulation in human macrophages. *Fitoterapia*. 2008; 79: 515-23.

80. Wannissorn B, Jarikasem S, Siriwanachai T, Thubthimthed S. Antibacterial properties of essential oils from Thai medicinal plants. *Fitoterapia*. 2005; 76: 233-6.

81. Ahmet AZ, Medine GLLCE, Meryem PENGL, Hatice UTC, Fikrettin PAHUN. Üsa K: Antimicrobial Effects of *Ocimum basilicum* (Labiatae) Extract. *Turk. J. Biol.* 2005; 29: 155-60.

82. Opalchenova G, Obreshkova D. Comparative studies on the activity of basil--an essential oil from *Ocimum basilicum* L.-against multidrug resistant clinical isolates of the genera *Staphylococcus*, *Enterococcus* and *Pseudomonas* by using different test methods. *J. Microbiol. Methods*. 2003; 54:105-10.

83. Patel VK, Bhatt VH. Folklore therapeutic indigenous plants in periodontal disorders in India (review, experimental and clinical approach). *Int. J. Clin. Pharmacol. Ther. Toxicol.* 1988; 26: 176-84.

84. Yamasaki K, Nakano M, Kawahata T, Mori H, Otake T, Ueba N, et al . Anti-HIV-1 activity of herbs in Labiateae. *Biol. Pharm. Bull.* 1998; 21: 829-33.

85. Renzulli C, Galvano F, Pierdomenico L, Speroni E, Guerra MC. Effects of rosmarinic acid against aflatoxin B1 and ochratoxin-A-induced cell damage in a human hepatoma cell line (Hep G2). *J. Appl. Toxicol.* 2004; 24: 289-96.

86. Manosroi J, Dhumtanom P, Manosroi A. Antiproliferative activity of essential oil extracted from Thai medicinal plants on KB and P388 cell lines. *CancerLett.* 2006; 235: 114-20.

87. Mohan L, Amberkar MV, Kumari M. *Ocimumsanum* L. (Tulsi) - An Overview — A Review Article. *Internat. J. Pharmaceu. Sci. — Rev and Resear* 2011; 7: 51-3.

88. Sandeep Pandey, Phytochemical Constituents, Pharmacological and Traditional Uses of *Ocimum gratissimum* L in Tropics, *Indo Am. J. P. Sci*, 2017; 4(11).

89. Iqbal J, Mishra RP. In vitro activity of medicinaplants against some bacterial and fungaisolates. *Asian J Pharm Clin Res* 2015;8:225-30.

90. Aziba PI, Bass D, Elegbe Y. Pharmacological investigation of *Ocimum gratissimum* in rodents *Phytother Res* 1999;13:427-29.

91. Iwalokun BA, Gbenle GO, Adewole TA, SmithSI, Akinsinde KA, Omonigbehin EO. Effects of *Ocimum gratissimum* L essential oil at subinhibitory concentrations on virulent and multidrug-resistant *Shigella* strains from Lagos, Nigeria. *APMIS* 2003;111:477-82.

92. Ekwensi MM, Oluigbo J, Akpuaka AAntibacterial activity of n-hexane extract of *Ocimum gratissimum* leaves. *IOSR J App Chem* 2014;7:6-10. 49.

93. Mann A. Phytochemical Constituents and Antimicrobial and Grain Protectant Activities of Clove Basil (*Ocimum gratissimum* L.) Grown in Nigeria. *Int J Plant Res* 2012;2:51-58.

94. Nweze EI, Eze E. Justification for the useof *Ocimum gratissimum* L in herbal medicine and its interaction with disc antibiotics. *BMC Compl Alter Med* 2009;9.

95. Ilori M, Sheteolu AO, Omonibgehin EA, Adeneye AA. Antidiarrhoeal activities of *Ocimum gratissimum* (Lamiaceae). *J Diarrhoeal Dis Res* 1996;14:283-85.

96. Pratheeba T, Ragavendran C, Natarajan D. Larvicidal, pupicidal and adulticidal potential of *Ocimum gratissimum* plant leaf extracts against filariasis inducing vector. *Int J Mosquito Res* 2015;2:1-8.

97. Macdonald IO, Oludare AS, Olabiyi A. Phytotoxic and Anti-microbial activities of Flavonoids in *Ocimum gratissimum*. *Life Sci J* 2010;7:45-48.

98. Mahapatra SK, Chakraborty SP, Das S, Roy S. Methanol extract of *Ocimum gratissimum* protects murine peritoneal macrophages from nicotine toxicityby decreasing free radical generation, lipid andprotein damage and enhances antioxidant protection. *Oxi Med Cell Long* 2009;2:222-30.

99. Nwanjo HU, Oze GO. Hypolipidaemic and Antioxidant properties of *Ocimum gratissimum* on Diabetic rats. *Plant Prod Res J* 2007;11:1-4.

100. Egesie UG, Adelaiye AB, Ibu JO, Egesie OJ. Safety and hypoglycaemic properties of aqueous leaf extract of *Ocimum gratissimum* in streptozotocininduced diabetic rats. *Nigerian J Physiol Sc* 2006;21:31-35.

101. Nangia-Makker P et al. Inhibition of breast tumor growth and angiogenesis by a medicinal herb: *Ocimum gratissimum*. *Int J Cancer* 2007;121:884-94.
102. Iweala EEJ, Liu f, Cheng R, Li Y, Omonhinmin CA, Zhang Y. Anticancer and free radical scavenging activity of some Nigerian food plants in vitro. *Int J Cancer Res* 2015;11:41-51.
103. Thomas N et al. Methanolic crude leaf extract of *Ocimum gratissimum* reverses phenylhydrazine-induced anemia in albino wistar rats. *Niger J Exp Clin Bio sci* 2013;1:23-7.
104. Ofem OE, Ani EJ, Eno AE. Effect of aqueous leaves extract of *Ocimum gratissimum* on hematological parameters in rats. *Int J Appl Basic Med Res* 2012;2:38-42.
105. Obianime AW, Aprioku JS, Esomonu C. The effects of aqueous *Ocimum gratissimum* leaf extract on some biochemical and hematological parameters in male mice. *Asian J Biol Sci* 2011;4:44-52.
106. Tripathi RD, Banerji R, Sharma ML, Balasubrahmanyam VR, Nigam SK. Toxicity of essential oil from a new strain of *Ocimum gratissimum* (Clocimum) against betel vine pathogenic fungi. *Agri Bio Chem* 1985;49:2277-82.
107. Hakkim FL, Arivazhagan G, Boopathy R. Antioxidant property of selected *Ocimum* species and their secondary metabolite content, *Journal of Medicinal Plants Research*, 2008, 2(9), 250-257.
108. Mwangi PW, Wambugu S, Kariuki DK, Mbugua PM, Kanui TI. Antinociceptive activities of the ethanolic extracts of *Ocimum kilimandscharicum* baker ex Gürke and *Ocimum kenyense* ayob. Ex a.j. Paton leaves. *IJP*. 2012; 3(1): 1-4..
109. Owuor BO, Ochanda JO, Kokwaro JO, Cheruiyot AC, Yeda RA, Okudo CA, et al. In vitro antiplasmoidal activity of selected Luo and Kuria medicinal plants. *J Ethnopharmacol*. 2012; 144:779-81.
110. Runyoro DKB, Ngassapa O, Kachali L, Obare V, Lyamuya EF. Biological activities of essential oils from plants growing in Tanzania. *East and Central African Journal of Pharmaceutical Sciences* 2010; 12: 85-91.
111. Sarin RV, Narwal S, Bafna PA. Anti-diarrheal activity of aqueous extract of *Ocimum kilimandscharicum*. *J Ethnopharmacol*. 2013; 148: 223-8.
112. Kumar AA, Mohan M, Haider SZ, Sharma A. Essential oil composition and antimicrobial activity of three *Ocimum* species from uttarakhand (India), *International journal of pharmacy and pharmaceutical sciences*, 2011, 3, 223-225.
113. Shinde K, Shinde V, Mahadik K, Gibbons S. Phytochemical and antibacterial studies on *Ocimum kilimandscharicum*. *Planta Med* 2010; 76: 412.
114. Monga J, Sharma M, Tailor N, Ganesh N. Antimelanoma and radioprotective activity of alcoholicaqueous extract of different species of *Ocimum* in C57BL mice. *Pharm Biol* 2011; 49(4): 428-36.
115. Jembere B, Obeng-Ofori D, Hassanali A. Products derived from the leaves of *Ocimum kilimandscharicum* Gürke (Labiatae) as post-harvest grain protectant against the infestation of three major store product insect pests, *Bulletin of Entomological Research*, 1995, 85,361-367.
116. Mahesh SP, Patil M B, Kumar R, Patil SR, Evaluation of aqueous extract of leaves of *Ocimum kilimandscharicum* Gürke wound healing activity in albino wistar rats, *International journal of pharmtech research coden (USA)*, 2009, 1, 544-550.
117. Rahman MS, Khan MMH, Jamal MA. Antibacterial. Anti-bacterial evaluation and minimum inhibitory concentration analysis of *Oxalis corniculata* and *Ocimum sanctum* against bacterial pathogens. *Biotechnolology*. 2010;9:533-6.
118. Farivar TN, Fard AHM, Zahedani SS, Naderi M, Moud BS. Anti tuberculosis effect of *Ocimum sanctum* extracts in in-vitro and macrophage culture. *J Medicinal Sci*. 2006;6:348-51.
119. Siva M, Shanmugam KR, Shanmugam B, Venkata SG, Ravi S, Sathyavelu RK, Mallikarjuna K. *Ocimum sanctum*: a review on the pharmacological properties. *Int J Basic Clin Pharmacol* 2016;5:558-65.
120. Khan A, Ahmad A, Manzoor N, Khan LA. Antifungal activities of *Ocimum sanctum* essential oil and its lead molecules. *Nat Prod Commun*. 2010;5(2):345-9.
121. Kelm MA, Nair MG, Strasburg GM, DeWitt DL. Antioxidant and cyclooxygenase inhibitory

phenolic compounds from *Ocimum sanctum* Linn. *Phytomedicine*. 2000;7:7-13.

122. Geetha RK, Vasudevan DM. Inhibition of lipid peroxidation by botanical extracts of *Ocimum sanctum*: In-vivo and in-vitro studies. *Life Sci.* 2004;76:21-8.

123. Siva M, Shanmugam KR, Shanmugam B, Venkata SG, Ravi S, Sathyavelu RK, Mallikarjuna K. *Ocimum sanctum*: a review on the pharmacological properties. *Int J Basic Clin Pharmacol* 2016;5:558-65.

124. Aruna K, Sivaramakrishnan VM. Anticarcinogenic effects of some Indian plant products. *Food Chem Toxicol.* 1992;30:953-6.

125. Ganasoundari A, Umadevi P, Rao BSS. Enhancement of bone marrow radioprotection and reduction of WR-2721 toxicity by *Ocimum sanctum*. *Mutation Res.* 1998;397:303-12.

126. Shetty SS, Udupu S, Udupu A. Evaluation of antioxidant and wound healing effects of alcoholic and aqueous extract of *Ocimum sanctum* Linn. In rats evid based complement. *Alternat Med.* 2008;5:95-110.

127. Govind P, Madhuri S. Medicinal plants: better remedy for neoplasm. *Indian Drug.* 2006;43(11):869-74.

128. Mediratta PK, Dewan V, Bhattacharya SK, Gupta VS, Maiti S, Sen P. Effect of *Ocimum sanctum* Linn. On humoral immune responses. *Indian J Med Res.* 1998;87:384.

129. Godhwani S, Godhwani JL, Vyas DS. *Ocimum sanctum* a preliminary study evaluating its immunoregulatory profile in albino rats. *J Ethnopharmacol.* 1988;24:193-8.

130. Sridevi G, Gopkumar P, Ashok S, Shastry CS. Pharmacological basis for antianaphylactic, antihistaminic and mast cell stabilization activity of *Ocimum sanctum*. *Inetrnet J Pharmacol.* 2009;7.

131. Ravindran R, Devi RS, Samson J, Senthilvelan M. Noise stress induced brain neurotransmitter changes and the effect of *Ocimum sanctum* (Linn) treatment in albino rats. *J Pharmcol Sci.* 2005;98:354-60.

132. Asha MK, Prashanth D, Murali B, Padmaja R, Amit A. Anthelmintic activity of essential oil of *Ocimum sanctum* and eugenol. *Fitoterapia.* 2001;72(6):66970.

133. Jaggi RK, Madaan R, Singh B. Anticonvulsant potential of holy basil, *Ocimum sanctum* Linn. And its cultures. *Indian J Exp Biol.* 2003;41:1329-33

134. Sood S, Narang D, Dinda AK, Maulik SK. Chronic oral administration of *Ocimum sanctum* Linn augments cardiac endogenous antioxidants and prevents isoproterenol-induced myocardial necrosis in rats. *J Pharm Pharmacol.* 2010;57:127-33.

135. Joshi H, Parle M. Evaluation of nootropic potential of *Ocimum sanctum* Linn. In mice. *Indian J Exp Biol.* 2006;44:133-6.

136. Komal S, Verma RJ. Protection against butyl hydroxybenzoic acid induced oxidative stress by *Ocimum sanctum* extract in mice liver. *Acta Poloniae Pharmaceutica Drug Resear* 2012;69(5):865-70.